TM1e

EoS Submission Details

EoS name	TM1e
category	Inner crust
submitted by	Helena Pais
affiliation	University of Coimbra
e-mail contact	hpais@uc.pt
sheet creation date	April 27, 2023

Abstract

This inner crust EoS, including nonspherical pasta phases, was calculated within a selfconsistent Thomas-Fermi approach [1] for β -equilibrium matter at zero temperature. The EoS parameterisation used was the non-linear RMF TM1e model [2]. This inner crust EoS has been used in Ref. [3]. Matched to model BsK22 [4] for the description of the outer crust below $n_B = 2 \times 10^{-3}$ fm⁻³.

References to the original work

- F. Grill. H. Pais, C. Providência, I. Vidaña, and S. Avancini, Phys. Rev. C 90, 045803 (2014).
- 2. H. Shen, F. Ji, J. Hu, and K. Sumiyoshi, Astrophys. J. 891, 148 (2020).
- O. Boukari, H. Pais, S. Antić, and C. Providência, Phys. Rev. C 103, 055804 (2021).
- J. M. Pearson, N. Chamel, A. Y. Potekhin, A. F. Fantina, C. Ducoin, A. K. Dutta, and S. Goriely, MNRAS 481, 2994 (2018); https://compose.obspm.fr/eos/210

Nuclear Matter Properties¹

	Quantity	Unit	
n_S	saturation density in symmetric matter	fm^{-3}	0.145
E_0	binding energy per baryon at saturation	MeV	-16.3
K	incompressibility	MeV	281
K'	skewness	MeV	-285
J	symmetry energy	MeV	31.4
L	symmetry energy slope parameter	MeV	40
K_{sym}	symmetry incompressibility	MeV	3.57

Neutron Star Properties¹

	Quantity	Unit	
M_{max}	maximum mass	M_{sun}	2.12
$M_{DU,e}$	mass at DUrca threshold (1/9) w/o μ^-	M_{sun}	0
$R_{M_{max}}$	radius at maximum NS mass	km	11.88
$R_{1.4}$	radius at $1.4 M_{sun} NS mass$	km	13.16
$ ilde{\Lambda}$	tidal deformability for GW170817 at a mass ratio of $q = 0.8$	-	770.15

eos.thermo

eos.thermo and the three grid defining files are CompOSE standard data files and by definition available. In eos.thermo, an extra quantity is added (last column). It refers to the geometry of the correspondent pasta phase, represented by an integer, with 0 for the outer crust, and 1 - droplets, 2 - rods, 3 - slabs, 4- tubes, 5- bubbles, and 6 corresponds to the core (homogeneous matter).

table dimension1table type1total number of grid points303

Range and density (#) of the grid parameters:

	Quantity	Unit	min	\max	#	
Т	Temperature	MeV	0	0	1	
\mathbf{n}_b	Baryon Nr Density	${\rm fm}^{-3}$	0.467962×10^{-9}	$0.89 imes 10^{-1}$	303	
\mathbf{Y}_q	Charge Fraction		0	0	1	

T, n_b, and Y_q are stored in eos.t, eos.nb, and eos.yq, respectively.

¹0-values indicate that the corresponding data is not provided.

Further Available Data Files

Files and quantities listed in the following are provided beyond CompOSE's core requirements as outlined in Sec.4.2. of the CompOSE manual.

eos.mr : This file provides the gravitational mass (in solar masses), the radius (in km), and the tidal deformability of a family of stars computed for this unified innercrust-core RMF EoS model, with the BsK22 outer crust.

eos.compo : available

4 particle pairs (neutrons, protons, electrons, muons) and one quadruple for heavy nucleus.

Phase index # 0: outer crust

Phase index # 7: inner crust

Phase index # 6: homogeneous matter

eos.micro : not available