

PT(GRDF2-DD2) general purpose EoS

EoS Submission Details

| | |
|---------------------|-----------------------------------------------------------|
| EoS name | PT(GRDF2-DD2) general purpose EoS |
| category | hadronic |
| submitted by | Stefan Typel |
| affiliation | Technische Universität Darmstadt, Institut für Kernphysik |
| e-mail contact | stypel@ikp.tu-darmstadt.de |
| sheet creation date | April 27, 2023 |

Abstract

This general-purpose hadronic EoS table is calculated with a generalized relativistic density functional (GRDF2) based on a relativistic mean-field model of nuclear matter with density dependent nucleon-meson couplings using the functional dependence introduced in [5.] and the parametrization DD2 [1.]. The dissolution of clusters and nuclei is described with medium-dependent mass shifts as defined in [3.]. Two-nucleon correlations in the continuum are considered as effective resonances with medium dependent properties [4.]. See [1.,2.,3.,6.] for details of the EoS model. Besides nucleons, electrons and muons with experimental masses, photons and nuclei are included as degrees of freedom. Masses of nuclei are taken from the 2016 Atomic mass evaluation [7.] if available. For other nuclei the predictions of the DZ31 model [8.] are used. The table entries give the properties of the 'local' EoS, i.e., the results of the density functional for the given temperature, density and charge fraction, without a phase transition construction.

References to the original work

1. S. Typel, G. Röpke, T. Kläahn, D. Blaschke, and H. H. Wolter,
Phys. Rev. C 81 (2010) 015803.
doi:10.1103/PhysRevC.81.015803
2. S. Typel,
J. Phys. G 45 (2018) 11400.
doi:10.1088/1361-6471/aadea5
3. H. Pais and S. Typel,
Comparison of equation of state models with different cluster dissolution mechanisms, in Nuclear Particle Correlations and Cluster Physics, edited by W. U. Schröder (World Scientific, Singapore) 2017, pp. 95-132.
doi:10.1142/9789813209350_0004

4. M. D. Voskresenskaya and S. Typel,
Nucl. Phys. A 887 (2012) 42.
doi:10.1016/j.nuclphysa.2012.05.006

Further References

5. S. Typel and H. H. Wolter,
Nucl. Phys. A 656 (1999) 331.
doi:10.1016/S0375-9474(99)00310-3
6. S. Typel S., H. H. Wolter, G. Röpke, and D. Blaschke,
Eur. Phys. J. A 50 (2014) 17.
doi:10.1140/epja/i2014-14017-x
7. Wang Meng, G. Audi, F. G. Kondev, W. J. Huang , S. Naimi and Xu Xing,
Chinese Physics C 41 (2017) 030003.
doi:10.1088/1674-1137/41/3/030003
8. J. Duflo and A. P. Zuker,
Phys. Rev. C 52 (1995) R23.
doi:10.1103/PhysRevC.52.R23

Nuclear Matter Properties

| | Quantity | | Unit | |
|-----------|-----------------------------------------|------------------|----------|--|
| n_S | saturation density in symmetric matter | fm^{-3} | 0.149065 | |
| E_0 | binding energy per baryon at saturation | MeV | 16.02 | |
| K | incompressibility | MeV | 242.72 | |
| K' | skewness | MeV | 168.65 | |
| J | symmetry energy | MeV | 32.67 | |
| L | symmetry energy slope parameter | MeV | 55.04 | |
| K_{sym} | symmetry incompressibility | MeV | -93.23 | |

Neutron Star Properties

| | Quantity | | Unit | |
|-------------------|---------------------------------------------------------------|-----|-----------|-------|
| M_{max} | maximum mass | | M_{sun} | 2.42 |
| $M_{DU,e}$ | mass at DURca threshold (1/9) w/o μ^- | | M_{sun} | 1.95 |
| $R_{M_{max}}$ | radius at maximum NS mass | | km | 11.87 |
| $R_{1.4}$ | radius at 1.4 M_{sun} NS mass | | km | 13.17 |
| $\tilde{\Lambda}$ | tidal deformability for GW170817 at a mass ratio of $q = 0.8$ | 789 | | |

eos.thermo

eos.thermo and the three grid defining files are CompOSE standard data files and by definition available.

No additional quantities are defined in eos.thermo.

| | |
|-----------------------------|---------|
| table dimension | 3 |
| table type | 1 |
| total number of grid points | 1830080 |

Range and density (#) of the grid parameters:

| Quantity | Unit | min | max | # |
|-------------------------|------------------|------------|-------|-----|
| T Temperature | MeV | 0.1 | 100 | 76 |
| n_b Baryon Nr Density | fm^{-3} | 10^{-10} | 1.000 | 301 |
| Y_q Charge Fraction | | 0.01 | 0.80 | 80 |

T, n_b , and Y_q are stored in eos.t, eos.nb, and eos.yq, respectively.

Further Available Data Files

Files and quantities listed in the following are provided beyond CompOSE's core requirements as outlined in Sec.4.2. of the CompOSE manual.

eos.compo : available

| index | particle |
|------------------|-----------------------------------------|
| 0 | e^- |
| 1 | μ^- |
| 10 | n |
| 11 | p |
| 700 | nn (1S_0) continuum correlations |
| 701 | np (1S_0) continuum correlations |
| 702 | pp (1S_0) continuum correlations |
| 2001 | ${}^2\text{H}$ |
| 3001 | ${}^3\text{H}$ |
| 3002 | ${}^3\text{He}$ |
| 4002 | ${}^4\text{H}$ |
| - end of table - | |

The tabulated particle number fractions of nucleons and leptons are net quantities, i.e., they are given by the difference of the corresponding particle and anti-particle fractions. A further particle set is defined. One set of quadruples gives the properties of a average heavy nucleus.

| index | particle |
|-------|-------------------|
| 0 | set of all nuclei |
| | - end of table - |

eos.micro : available

| index | quantity | particle |
|-------|---------------------------------------|----------|
| 10051 | relativistic vector self-energy V_n | n |
| 10052 | relativistic scalar self-energy S_n | n |
| 11051 | relativistic vector self-energy V_p | p |

- continued on next page -

| | | |
|-------|---------------------------------------|-----------------------|
| index | quantity | particle |
| 11052 | relativistic scalar self-energy S_p | p - end of table - |

Description of Phases

The table contains the locally calculated EoS data without performing a phase transition construction.

| | | |
|-------------|--------------------------|--|
| phase index | description | |
| 0 | mixture of all particles | |